

# Enhanced exercise capacity in mice with severe heart failure treated with an allosteric effector of hemoglobin, *myo*-inositol trispyrophosphate

Andreia Biolo<sup>a</sup>, Ruth Greferath<sup>b,c</sup>, Deborah A. Siwik<sup>a</sup>, Fuzhong Qin<sup>a</sup>, Eugene Valsky<sup>a</sup>, Konstantina C. Fylaktakidou<sup>c,1</sup>, Srinivasu Pothukanuri<sup>c</sup>, Carolina D. Duarte<sup>c</sup>, Richard P. Schwarz<sup>b</sup>, Jean-Marie Lehn<sup>c,2</sup>, Claude Nicolau<sup>b,c,d,2</sup>, and Wilson S. Colucci<sup>a,2</sup>

<sup>a</sup>Cardiovascular Section, Department of Medicine, Boston University Medical Center, and Myocardial Biology Unit, Boston University School of Medicine, Boston, MA 02115; <sup>b</sup>NormOxys, Inc., 200 Boston Avenue, Medford, MA 02155; <sup>c</sup>Institut de Science et d'Ingénierie Supramoléculaires, Université Louis Pasteur, 8 Allée Gaspard Monge, 67000 Strasbourg, France; and <sup>d</sup>Friedman School of Nutrition Science and Policy, Tufts University, Boston, MA 02115

Contributed by Jean-Marie Lehn, December 31, 2008 (sent for review December 1, 2008)

A major determinant of maximal exercise capacity is the delivery of oxygen to exercising muscles. *myo*-inositol trispyrophosphate (ITPP) is a recently identified membrane-permeant molecule that causes allosteric regulation of Hb oxygen binding affinity. In normal mice, i.p. administration of ITPP (0.5–3 g/kg) caused a dose-related increase in the oxygen tension at which Hb is 50% saturated (p50), with a maximal increase of 31%. In parallel experiments, ITPP caused a dose-related increase in maximal exercise capacity, with a maximal increase of  $57 \pm 13\%$  ( $P = 0.002$ ). In transgenic mice with severe heart failure caused by cardiac-specific overexpression of  $G\alpha_q$ , i.p. ITPP increased exercise capacity, with a maximal increase of  $63 \pm 7\%$  ( $P = 0.005$ ). Oral administration of ITPP in drinking water increased Hb p50 and maximal exercise capacity ( $+34 \pm 10\%$ ;  $P < 0.002$ ) in normal and failing mice. Consistent with increased tissue oxygen availability, ITPP decreased hypoxia inducible factor-1 $\alpha$  mRNA expression in myocardium. It had no effect on myocardial contractility in isolated mouse cardiac myocytes and did not affect arterial blood pressure in vivo in mice. Thus, ITPP decreases the oxygen binding affinity of Hb, increases tissue oxygen delivery, and increases maximal exercise capacity in normal mice and mice with severe heart failure. ITPP is thus an attractive candidate for the therapy of patients with reduced exercise capacity caused by heart failure.

hypoxia | oxygen delivery

A major determinant of exercise capacity is the amount of oxygen available to the exercising muscles. Heart failure is characterized by a reduction in exercise capacity caused primarily to the inability of the cardiovascular system to increase cardiac output and hence, blood flow, to exercising muscles (1–5). Accordingly, most previous efforts to improve exercise capacity in patients with heart failure have relied on interventions that increase the pumping function of the heart and/or the distribution of blood flow to exercising muscles. Increasing the concentration of Hb also enhances the delivery of oxygen to exercising muscles, and in patients with concurrent heart failure and anemia the administration of erythropoietin to increase the red cell mass may increase maximal exercise capacity (6, 7).

The increased delivery of oxygen to exercising muscles might also be achieved at a constant Hb concentration by decreasing the oxygen binding affinity of Hb so as to release more oxygen in hypoxic tissues. The organic phosphate 2,3-bisphosphoglycerate (2,3-BPG) is the natural allosteric effector that decreases the oxygen binding affinity of human Hb, and increased levels of 2,3-BPG appear to play a compensatory role in a variety of circumstances including high altitude and chronic pulmonary disease (8, 9). It has been noted that red blood cell 2,3-BPG levels are increased in patients with low output heart failure, leading to the suggestion that interventions to further decrease

Hb oxygen binding affinity might be of clinical value in such patients (10).

*myo*-inositol hexakisphosphate (IP<sub>6</sub>) is a powerful allosteric effector of Hb that increases the regulated oxygen-releasing capacity of RBC. In piglets, transfusion of RBC loaded ex vivo with IP<sub>6</sub> led to physiologic effects consistent with increased oxygen delivery (11, 12). Because at neutral pH, IP<sub>6</sub> is a partially dissociated polyanion bearing 8 negative charges (13), it is unable to cross the RBC membrane and ex vivo physical methods are required to load erythrocytes (11, 14). However, we recently showed that *myo*-inositol trispyrophosphate (ITPP) hexasodium salt is capable of crossing the RBC plasma membrane and acting as an allosteric effector of Hb, shifting the oxyhemoglobin dissociation curve to higher oxygen pressures (pO<sub>2</sub>) (15, 16), and leading to a marked inhibition of blood vessel formation processes in vitro (16). Accordingly, we tested the hypothesis that systemic administration of ITPP would increase exercise capacity in normal mice and in mice with severe exercise limitation caused by reduced cardiac output as a result of myocardial failure.

## Results and Discussion


**ITPP Decreases the Oxygen Binding Affinity of Hb in Vivo.** To determine whether ITPP increases the tissue availability of oxygen in vivo, ITPP was administered to normal mice by i.p. injection and the oxygen binding affinity of Hb was measured 18–24 h later. In blood from control mice, the pO<sub>2</sub> at which Hb is 50% saturated with oxygen (p50) averaged  $37.3 \pm 1.5$  torr. ITPP caused a dose-related right shift in the oxygen dissociation curve, resulting in a mean 22% increase in p50 at a dose of 1 g/kg, and a mean 37% increase with the 2 g/kg dose (Fig. 1A). There was little or no effect of ITPP on maximal oxygen saturation at pO<sub>2</sub> of  $>100$  torr (Fig. 1B). Thus, although ITPP had no significant effect on total oxygen carrying capacity, the right shift in the oxygen dissociation curve at O<sub>2</sub> tensions in the physiologic range suggests that a larger fraction of O<sub>2</sub> would be released at the lower oxygen tension in the tissue. After a single ITPP administration, the p50 increase was sustained for 48 h, had decreased by  $\approx 50\%$  in 5 days, and was no longer present after 12 days. The prolonged duration of the effect, which is consistent with the

Author contributions: J.-M.L., C.N., and W.S.C. designed research; and A.B., R.G., D.A.S., F.Q., E.V., K.C.F., S.P., C.D.D., and R.P.S. performed research.

Conflict of interest statement: W.S.C., R.G., J.-M.L., and C.N. own stock in NormOxys, Inc., which holds the patents on the applications of inositol trispyrophosphate. R.G., J.-M.L., and C.N. are listed as coinventors together with K.C.F. on these patents. In addition, R.G., C.N., and S.P. are employed by NormOxys, Inc.

<sup>1</sup>Present address: Molecular Biology and Genetics Department, Democritus University of Thrace, Dimitris 19, 68100 Alexandroupolis, Greece.

<sup>2</sup>To whom correspondence may be addressed. E-mail: lehn@isis.u-strasbg.fr, claude.nicolau@normoxys.com, or wilson.colucci@bmc.org.



It has recently been reported that, in mice, bone marrow transplantation with a Hb variant having a low oxygen affinity or

PNAS | February 10, 2009 | vol. 106 | no. 6 | 1927

i.v. administration of another type of allosteric Hb modifier increased the voluntary running time (19). Whereas measurement of voluntary running time precludes conclusions about possible effects on maximal exercise capacity, we used here an exercise protocol designed to measure maximal exercise capacity, which is directly related to the maximal oxygen delivery.

Our findings suggest that increasing the availability of oxygen with ITTP may be a means of improving maximal exercise in patients with heart failure who have impaired exercise capacity caused by reduced cardiac output. However, it should be noted that the causes of reduced exercise in patients with heart failure are incompletely understood and may involve multiple factors other than oxygen delivery, including deconditioning, atrophy, and/or metabolic dysfunction of the skeletal (18). It is also possible that the Gαq mouse model of dilated heart failure may not fully reflect these and other causes of exercise impairment in humans.

**Oral ITTP Increases p50 and Exercise Capacity.** Previous efforts to modify RBC Hb affinity pharmacologically have been limited by the failure of effector molecules to cross RBC membranes (11, 14), short duration of effects (19), and poor solubility in water (20). Because ITTP is both membrane permeant and readily soluble in water (15, 16), we considered the possibility that ITTP would be absorbed after oral ingestion.

To determine whether oral ingestion of ITTP causes a shift in Hb oxygen affinity, normal mice were allowed to drink ad libitum water in which ITTP or IP<sub>6</sub> was dissolved at a concentration of 20 mg/mL. Hb p50 increased by ≈16% in mice that drank ITTP, but was unchanged in mice that drank water (+3%) or water with IP<sub>6</sub> (+3%), which is membrane impermeant. To determine whether oral ingestion would also improve exercise function, maximal exercise capacity was determined in normal mice ( $n = 4$ ) and mice with heart failure caused by Gq overexpression ( $n = 2$ ) before and after drinking water containing ITTP (20 mg/mL, ad libitum) for 4–8 days. Oral ingestion of ITTP increased maximal exercise capacity by an average of  $34 \pm 10\%$  ( $P < 0.002$ ;  $n = 6$ ).

**Lack of Direct Vascular and Cardiac Effects of ITTP.** Pharmacologic agents that exert vasodilator and/or positive inotropic effects may increase cardiac output, blood flow to skeletal muscle, and exercise capacity in heart failure (21). To determine whether ITTP exerts vasodilator effects, blood pressure was measured in 12 normal mice at baseline and 24 h after i.p. injection of ITTP (2 g/kg) or placebo. Neither ITTP nor placebo affected systolic blood pressure (before ITTP =  $134 \pm 6$  mm Hg; after ITTP =  $135 \pm 4$  mm Hg;  $P$  not significant), diastolic blood pressure (before ITTP =  $76 \pm 12$  mm Hg; after ITTP =  $73 \pm 4$  mm Hg;  $P$  not significant), or heart rate (before ITTP =  $706 \pm 12$ ; after ITTP =  $720 \pm 17$  beats per min;  $P$  not significant).

To determine whether ITTP exerts a direct positive inotropic effect, contractile properties were measured in freshly-isolated adult rat ventricular myocytes, as described (22). ITTP (500 μM; 10 min) had no effect on baseline myocyte sarcomere shortening ( $-9 \pm 29\%$ ;  $P$  not significant) or on the rate of sarcomere shortening ( $-17 \pm 27\%$ ;  $P$  not significant), whereas the positive control norepinephrine caused 4- to 6-fold increases in sarcomere shortening ( $+405 \pm 155\%$ ,  $P < 0.01$ ) and the rate of shortening ( $+670 \pm 170\%$ ;  $P < 0.01$ ).

## Conclusion

The present results show that i.p. and oral administration of ITTP cause a decrease in Hb-oxygen affinity that is associated

with a parallel, dose-related increase in maximal exercise capacity in both normal mice and mice with reduced exercise capacity caused by severe myocardial failure. These observations demonstrate that allosteric modulation of Hb oxygen binding affinity can exert clinically meaningful effects on maximal exercise capacity. Accordingly, ITTP is an attractive therapeutic candidate to alleviate symptoms in patients with reduced exercise capacity caused by low cardiac output heart failure. ITTP may also enhance physical performance of otherwise healthy individuals, in particular, under extreme conditions such as high altitude or intense physical exercise.

## Methods

**Synthesis of ITTP.** This membrane-permeant allosteric effector of Hb was synthesized as reported (15), by using an improved procedure. Briefly, IP<sub>6</sub> dodecasodium salt (Sigma) was converted via its perprotonated form, obtained by passage over Dowex 50 resin (Sigma-Aldrich) in its H<sup>+</sup> form, into its triethylammonium salt and then, upon heating with dicyclohexylcarbodiimide (Sigma-Aldrich) in acetonitrile/water 2/1 solution, triply cyclized to give the triethylammonium salt of ITTP. Thereafter, an aqueous solution of the latter was passed over Dowex Marathon C Na<sup>+</sup> resin (Sigma-Aldrich) until cation exchange of triethylammonium against sodium was complete. Evaporation of the aqueous solution gave the sodium salt of ITTP in high purity (≈90% yield).

**p50 Measurements.** The p50 value (pO<sub>2</sub> at which 50% of Hb is saturated with O<sub>2</sub>), a measure of the affinity of Hb for oxygen, was determined by using a HEMOX Analyzer (TCS Scientific), by constructing Hb dissociation curves based on dual wavelength spectrophotometry as described (16).

**RT-PCR for HIF-1α mRNA.** Total RNA was extracted from mouse hearts by using a Total RNA Purification System (Invitrogen). Quantitative RT-PCR was performed by using cybergreen and the I-cycler iQ RT-PCR (Bio-Rad). The primers for HIF-1α were: 5'-TCAAGTCAGCAACGTGGAAG-3' and 5'-TATCGAGGCTGTGTCGACTG-3'.

**Exercise Testing.** Maximum exercise capacity was measured by using a rodent treadmill equipped with an air puff motivator (Columbus Instruments) as described (23). Animals were familiarized with running on the treadmill. For tests, the treadmill was set at a constant incline of 15°, and the initial speed of 15 m/min was increased by 3 m/min every 2 min. Total exercise time was recorded as the elapsed time to exhaustion and then converted to distance. Exhaustion was determined by an observer blinded to treatment group and was defined as the point at which the animals could not keep pace with the treadmill and had no response to the air puff stimulus. All exercise evaluations were performed twice and the results were averaged.

**Gαq Transgenic Mice.** Gαq transgenic mice were kindly provided by Gerald W. Dorn II (University of Cincinnati, Cincinnati) (17) and subsequently bred at Boston University Medical Center. This mouse overexpresses Gαq exclusively in the myocardium under the myosin heavy chain promoter. The present study used the Gαq line that overexpresses 40 (Gαq40) copies of the transgene on a FVB background, which leads to a 5-fold increase in the transgene protein levels. Age-matched controls were bred by using heterozygote Gαq male mice with WT females obtained from Charles River. Animals were studied at 12–14 weeks of age. The Institutional Animal Care and Use Committee at Boston University School of Medicine approved all study procedures and use of animals.

**Statistical Analysis.** Data are shown as mean ± SEM. Group comparisons were performed by using Student's  $t$  test (2 groups) or 1-way ANOVA (multiple groups). Baseline versus treatment comparisons were performed by paired  $t$  test. All tests were 2-tailed, and  $P < 0.05$  was considered significant.

**ACKNOWLEDGMENTS.** We thank Dr. Elena Tufa for help with the p50 measurements. This work was supported by National Institutes of Health Grants HL61639 and HL20612 (to W.S.C.), Coordinating Agency for Higher Education (CAPES-BR) fellowships (to A.B. and C.D.D.), and a grant from NormOxys, Inc.

- Weber KT, Kinasewitz GT, Janicki JS, Fishman AP (1982) Oxygen utilization and ventilation during exercise in patients with chronic cardiac failure. *Circulation* 65:1213–1223.
- Sullivan MJ, Hawthorne MH (1995) Exercise intolerance in patients with chronic heart failure. *Prog Cardiovasc Dis* 38:1–22.

- Katz SD, et al. (2000) Near-maximal fractional oxygen extraction by active skeletal muscle in patients with chronic heart failure. *J Appl Physiol* 88:2138–2142.
- Pina IL, et al. (2003) Exercise and heart failure: A statement from the American Heart Association Committee on exercise, rehabilitation, and prevention. *Circulation* 107:1210–1225.

5. Givertz MM, Colucci WS, Braunwald E (2005) in *Braunwald's Heart Disease*, eds Zipes DP, Libby P, Bonow RO, Braunwald E (Elsevier Saunders, Philadelphia), pp 539–568.
6. Mancini DM, et al. (2003) Effect of erythropoietin on exercise capacity in patients with moderate to severe chronic heart failure. *Circulation* 107:294–299.
7. Palazzuoli A, et al. (2006) Erythropoietin improves anemia exercise tolerance and renal function and reduces B-type natriuretic peptide and hospitalization in patients with heart failure and anemia. *Am Heart J* 152:1096–1115.
8. Benesch R, Benesch RE, Yu CI (1968) Reciprocal binding of oxygen and diphosphoglycerate by human hemoglobin. *Proc Natl Acad Sci USA* 59:526–532.
9. Bunn HF, Jandl JH (1970) Control of hemoglobin function within the red cell. *N Engl J Med* 282:1414–1421.
10. Bersin RM, et al. (1993) Importance of oxygen-hemoglobin binding to oxygen transport in congestive heart failure. *Br Heart J* 70:443–447.
11. Teisseire BP, Ropars C, Vallez MO, Herigault RA, Nicolau C (1985) Physiological effects of high-P50 erythrocyte transfusion on piglets. *J Appl Physiol* 58:1810–1817.
12. Teisseire B, Ropars C, Villereal MC, Nicolau C (1987) Long-term physiological effects of enhanced O<sub>2</sub> release by inositol hexaphosphate-loaded erythrocytes. *Proc Natl Acad Sci USA* 84:6894–6898.
13. Poillon WN, Robinson MD, Kim BC (1985) Deoxygenated sickle hemoglobin. Modulation of its solubility by 2,3-diphosphoglycerate and other allosteric polyanions. *J Biol Chem* 260:13897–13900.
14. Bruggemann U, Roux EC, Hannig J, Nicolau C (1995) Low-oxygen-affinity red cells produced in a large-volume, continuous-flow electroporation system. *Transfusion* 35:478–486.
15. Fylaktakidou KC, Lehn J-M, Greferath R, Nicolau C (2005) Inositol tripyrophosphate: A new membrane permeant allosteric effector of hemoglobin. *Bioorg Med Chem Lett* 15:1605–1608.
16. Kieda C, et al. (2006) Suppression of hypoxia-induced HIF-1 $\alpha$  and of angiogenesis in endothelial cells by myo-inositol trispyrophosphate-treated erythrocytes. *Proc Natl Acad Sci USA* 103:15576–15581.
17. D'Angelo DD, et al. (1997) Transgenic G $\alpha_q$  overexpression induces cardiac contractile failure in mice. *Proc Natl Acad Sci USA* 94:8121–8126.
18. Minotti JR, Christoph I, Massie BM (1992) Skeletal muscle function, morphology, and metabolism in patients with congestive heart failure. *Chest* 101:3335–3395.
19. Watanabe T, et al. (2008) Reduction in hemoglobin-oxygen affinity results in the improvement of exercise capacity in mice with chronic heart failure. *J Am Coll Cardiol* 52:779–786.
20. Randad RS, Mahran MA, Mehanna AS, Abraham DJ (1991) Allosteric modifiers of hemoglobin. 1. Design, synthesis, testing, and structure-allosteric activity relationship of novel hemoglobin oxygen affinity decreasing agents. *J Med Chem* 34:752–757.
21. Clark AL (2006) Exercise and heart failure: Assessment and treatment. *Heart* 92:699–703.
22. Satoh N, Suter TM, Liao R, Colucci WS (2000) Chronic  $\alpha$ -adrenergic receptor stimulation modulates the contractile phenotype of cardiac myocytes in vitro. *Circulation* 102:2249–2254.
23. Trueblood NA, et al. (2005) Biphasic temporal pattern in exercise capacity after myocardial infarction in the rat: Relationship to left ventricular remodeling. *Am J Physiol* 288:H244–H249.